Recommendations for receiving the **MODERNA** Alternative COVID-19 Vaccine



The COVID-19 pandemic has still not ended, so the Ministry of Public Health recommends that after the third dose people continue to receive booster doses every 4 months to increase immunity.

First dose	Second T dose	Required interval after 2 nd dose	Third TTT dose	Required interval after 3 rd dose	Fourth Fourth dose
moderna	(4 weeks after first dose)	After 3 months	moderna	After 4 months	moderna
Sinovac /	(4 weeks after first dose)	After 3 months	moderna	After 4 months	moderna
AstraZeneca	(4 weeks after first dose)	After 3 months	moderna	After 4 months	moderna
Pfizer	(4 weeks after first dose)	After 3 months	moderna	After 4 months	moderna
Johnson	Afohnson	2 months	(Booster dose)	After 4 months	moderna
Sinovac /	SINOPHARM	After 4 weeks	moderna	After 4 months	moderna
Sinovac /	AstraZeneca	After 3 months	moderna	After 4 months	moderna
AstraZeneca	AstraZeneca	After 3 months	moderna	After 4 months	moderna
Sinovac /	Pfizer	After 3 months	moderna	After 4 months	moderna
Pfizer	Pfizer	After 3 months	moderna	After 4 months	moderna
AstraZeneca	Pfizer	After 3 months	moderna	After 4 months	moderna
Sinovac /	Sinovac /	After 4 weeks	AstraZeneca	After 4 months	moderna
Sinovac /	Sinovac /	After 4 weeks	Pfizer	After 4 months	moderna
Sinovac /	AstraZeneca	After 3 months	AstraZeneca	After 4 months	moderna
AstraZeneca	AstraZeneca	After 3 months	Pfizer	After 4 months	moderna

Note: If you have recovered from COVID-19, you can receive the COVID-19 vaccine 3 months after testing positive through the same recommendation as those who have never been infected with the COVID-19.

Children and teenagers aged 6-17 are urged to strictly follow the pediatrician's recommendations.